



LARIXIFOL Plant Health Supplement

An antioxidant supplement for plant health

LARIXIFOL is a highly purified taxifolin-based antioxidant derived from larch wood. When used with good plant nutritional practices, LARIXIFOL fortifies growing plants against stresses such as excess heat, cold, and drought.

ACTIVE INGREDIENT

Taxifolin* 12%
*bioflavonoid complex derived from larch wood (*Larix gmelini*)

GUARANTEED ANALYSIS

OTHER INGREDIENTS
propylene glycol 30%
soy lecithin 6%
water 52%

NET CONTENTS: 33.8 (Fluid Oz.) NET WEIGHT: 2.1 lb

DIRECTIONS FOR USE: Shake before opening. See following table for crops and rates:

CAUTION - KEEP OUT OF REACH OF CHILDREN

CROP	SEED TREATMENT APPLICATION	FOLIAR APPLICATION
Cucurbits (cucumbers, melons)	Mix 0.015 fl oz (1/10 teaspoon) of LARIXIFOL per 10 lb seed in 10 gallons water. Soak seed for 2 hours shortly before planting.	Mix 2.75 fl oz per acre in 40 - 50 gallons per acre. Apply as a low-pressure spray at the following stages: 1) the 2 - 3 leaf stage, 2) at flowering and 3) at the beginning of fruit set.
Small cereal grains (wheat, barley, rye, oats)	Mix 2.7 fl oz LARIXIFOL per ton of seed in 2.5 gallons water. Mix thoroughly with seed using appropriate equipment and allow to dry.	Mix 1 fl oz LARIXIFOL per acre in 80 gallons water per acre. Apply as a low pressure spray at the following stages: 1) Flag leaf stage (Feekes stage 7 - 8), 2) boot stage (Feekes stage 10)
Corn	Mix 3.1 fl. oz. LARIXIFOL in 2.6 gal water per ton of seed. Mix thoroughly with seed using appropriate equipment and allow to dry.	Mix 1.7 fl oz per acre in 32 gallons per acre. Apply as a low-pressure spray at the following stages: 1) the 3-5 leaf stage, and 2) the V6 stage of development
Grapes		Mix 3.4 fl oz LARIXIFOL per acre in 65 gallons water per acre. Apply as a low-pressure spray at the following intervals: 1) at flowering, 2) 20 days following first application.
Soybeans	Mix 3.1 fl oz per ton of seed in 2.5 gallons water per ton of seed. Mix thoroughly with seed using appropriate equipment and allow to dry.	Mix 1.1 fl oz LARIXIFOL per acre in 80 gallons water per acre. Apply as a low pressure spray at the following intervals: 1) initial flowering, 2) 14 days after first spray, 3) 30 days after second spray.
Peppers, Chiles		Mix 1.35 fl oz per 10 gallons water. Apply as a low pressure spray at the flowering stage.
Potatoes	Tuber treatment: Mix 0.6 fl oz LARIXIFOL per ton of tubers in 2.5 gallons water per ton of tubers. Mix thoroughly using appropriate equipment and allow to dry.	Mix 2.7 fl oz LARIXIFOL per acre in 80 gallons water per acre. Apply as a low pressure spray at the following intervals: 1) at budding, 2) 20 days after first spray
Tomatoes		Mix 0.8 fl oz LARIXIFOL per 5 acres tomatoes in 40 - 50 gallons water per acre. Apply as a low-pressure spray at the following intervals: - 1) initial flowering, 2) and 3) at 10 day intervals following first spray.
Sunflowers	Mix 3.1 fl oz LARIXIFOL per ton of seed in 2.5 gallons water per ton of seed. Mix thoroughly with seed using appropriate equipment and allow to dry.	Mix 2.7 fl oz LARIXIFOL per acre in 32 gallons water per acre. Apply as a low pressure spray at the following intervals: 1) Bud stage, 2) Flowering
Sugar beets	Mix 7.7 fl oz LARIXIFOL per ton of seed in 5 gallons water per ton of seed. Mix thoroughly with seed using appropriate equipment and allow to dry.	Mix 3.4 fl oz LARIXIFOL per acre in 80 gallons water per acre. Apply as a low-pressure spray at the following intervals: 1) 3- 4 leaf pairs, 2) beginning of canopy closing
Rice	Mix 3.1 fl. oz. LARIXIFOL in 2.5 gal water per ton of seed. Mix thoroughly with seed using appropriate equipment and allow to dry.	Mix 1.4 fl. oz. in 12 gallons per acre of water. Apply at during vegetative stage.

CONDITIONS OF SALE: The information contained on this label is believed to be accurate and reliable. Buyer and User acknowledge and assume all liability resulting from the use of this product. Follow directions carefully. Timing, method of application, weather, crop and soil conditions are beyond the control of the seller.

Manufactured for: AgroMarketing Company, Inc., 133 Mavey St., Toronto, ON Canada, M6P 2L8 Tel. 416-628-5147
Imported Exclusively by: AgroShield LLC, 914 Walnut Bend Lane, Houston TX 77042 Tel. 515-419-9524